


## AC115 *Hymenomonas coronata*

Taxonomy		Image	Origin		
<b>Kingdom</b>	Chromista		<b>Climatic Zone</b>	Subtropical	
<b>Division</b>	Haptophyta		<b>Habitat</b>	Coastal	
<b>Class</b>	Prymnesiophyceae		<b>Ocean</b>	-	
<b>Order</b>	Coccolithales		<b>Sea</b>	-	
<b>Family</b>	<i>Hymenomonadaceae</i>		<b>Country</b>	France	
<b>Genus</b>	<i>Hymenomonas</i>		<b>Region</b>	Guadeloupe	
<b>Species</b>	coronata		<b>Geolocation</b>		
<b>Authority(ies)</b>	-		<b>Latitude</b>	-	
		<b>Longitude</b>	-		
Identity		Morphology	Culture		
<b>Holotype</b>	No	<b>Length (µm)</b>	-	<b>Isolated by</b>	J. Fresnel
<b>Equivalent code</b>	-	<b>Width (µm)</b>	-	<b>Isolation date</b>	1984
<b>Deposited by</b>	-	<b>Diameter</b>	-	<b>Isolation site</b>	Guadeloupe
<b>Deposit date</b>	-	<b>Wall</b>	-	<b>Culture medium</b>	ES
<b>Identified by</b>	-	<b>Comments</b>	-	<b>Growth temperature</b>	16 °C
				<b>Growth light</b>	90 µE
				<b>Toxicity</b>	Unproven
External Links					
<b>Genbank accession number</b>	-				
<b>References</b>	<p>- Fresnel J. (1994) A heteromorphic life cycle in two coastal coccolithophorids, <i>Hymenomonas lacuna</i> and <i>Hymenomonas coronata</i> (Prymnesiophyceae). <i>Can. J. Bot.</i> 72: 1455-1462</p> <p>- Bittner L, Gobet A, Audic S, Romac S, Egge E.S., Santini S, Ogata H, Probert I, Edvardsen B and de Vargas C (2012) Diversity patterns of uncultured Haptophytes unravelled by pyrosequencing in Naples Bay. <i>Molecular Ecology</i> DOI: 10.1111/mec.12108</p>				

AC115 *Hymenomonas coronata*

