


## AC13 *Ochrosphaera neapolitana*

Taxonomy		Image		Origin	
<b>Kingdom</b>	Chromista			<b>Climatic Zone</b>	Mediterranean
<b>Division</b>	Haptophyta			<b>Habitat</b>	Coastal
<b>Class</b>	Prymnesiophyceae			<b>Ocean</b>	-
<b>Order</b>	Coccolithales			<b>Sea</b>	Mediterranean Sea
<b>Family</b>	<i>Hymenomonadaceae</i>			<b>Country</b>	France
<b>Genus</b>	<i>Ochrosphaera</i>			<b>Region</b>	Provence-Alpes-Côte d'Azur
<b>Species</b>	neapolitana			<b>Geolocation</b>	
<b>Authority(ies)</b>	Schussnig			<b>Latitude</b>	43.305
Identity		Morphology		Culture	
<b>Holotype</b>	No	<b>Length (µm)</b>	-	<b>Isolated by</b>	S. Maestrini
<b>Equivalent code</b>	-	<b>Width (µm)</b>	-	<b>Isolation date</b>	1967
<b>Deposited by</b>	-	<b>Diameter</b>	6,50 µm	<b>Isolation site</b>	Marseille
<b>Deposit date</b>	-	<b>Wall</b>	Coccosphere	<b>Culture medium</b>	ES
<b>Identified by</b>	-	<b>Comments</b>	CB: Observed forms: palmelloid stage with coccoliths; zoïdes (with coccoliths?) with two unequal flagella. CB 05/93: always in diploid phase according to JF but it seems like there are also haploid cells. Biovolume 0,144 pL	<b>Growth temperature</b>	16 °C
				<b>Growth light</b>	90 µE
				<b>Toxicity</b>	Unproven
External Links					
<b>Genbank accession number</b>	-				
<b>References</b>	- Bittner L, Gobet A, Audic S, Romac S, Egge E.S., Santini S, Ogata H, Probert I, Edvardsen B and de Vargas C (2012) Diversity patterns of uncultured Haptophytes unravelled by pyrosequencing in Naples Bay. <i>Molecular Ecology</i> DOI: 10.1111/mec.12108				

# AC13 *Ochrosphaera neapolitana*

