


AC14 *Sarcinochrysis marina*

Taxonomy		Image		Origin	
Kingdom	Chromista			Climatic Zone	Temperate
Division	Heterokontophyta			Habitat	Coastal
Class	Pelagophyceae			Ocean	North Atlantic
Order	Sarcinochrysidales			Sea	English Channel
Family	<i>Sarcinochrysidaceae</i>			Country	France
Genus	<i>Sarcinochrysis</i>			Region	Basse-Normandie
Species	marina			Geolocation	
Authority(ies)	Geitler			Latitude	49.319
Identity		Morphology		Culture	
Holotype	No	Length (µm)	6-7 µm	Isolated by	H. Lepailleur
Equivalent code	-	Width (µm)	4-5 µm	Isolation date	1965
Deposited by	-	Diameter	-	Isolation site	Luc sur Mer
Deposit date	-	Wall	-	Culture medium	ES
Identified by	-	Comments	Observed forms: palmelloid stage "sarcinochrysoïde"; zoïdes; old strains -> palmelloid stage similar to <i>Ruttnera chadefaudii</i> .	Growth temperature	16 °C
				Growth light	70 µE
				Toxicity	Unproven
External Links					
Genbank accession number	-				
References	- Gayral P. (1972). Sur les Chrysophycées à zoïdes pélophycéens, notamment <i>Sarcinochrysis marina</i> Geitler. Soc. Phycol de France, Bull 17: 40-45.				