


## AC223 *Nannochloropsis gaditana*

Taxonomy		Image		Origin	
<b>Kingdom</b>	Chromista			<b>Climatic Zone</b>	Temperate
<b>Division</b>	Heterokontophyta			<b>Habitat</b>	Coastal
<b>Class</b>	Eustigmatophyceae			<b>Ocean</b>	-
<b>Order</b>	Eustigmatales			<b>Sea</b>	-
<b>Family</b>	<i>Monodopsidaceae</i>			<b>Country</b>	Maroc
<b>Genus</b>	<i>Nannochloropsis</i>			<b>Region</b>	-
<b>Species</b>	<i>gaditana</i>			<b>Geolocation</b>	
<b>Authority(ies)</b>	Lubiàn			<b>Latitude</b>	32.739
				<b>Longitude</b>	-9.040000
Identity		Morphology		Culture	
<b>Holotype</b>	No	<b>Length (μm)</b>	4.09 μm	<b>Isolated by</b>	C. Billard
<b>Equivalent code</b>	EUS1	<b>Width (μm)</b>	-	<b>Isolation date</b>	juin 1985
<b>Deposited by</b>	-	<b>Diameter</b>	-	<b>Isolation site</b>	Oualidia
<b>Deposit date</b>	-	<b>Wall</b>	-	<b>Culture medium</b>	f/2
<b>Identified by</b>	C. Billard	<b>Comments</b>	-	<b>Growth temperature</b>	16 °C
				<b>Growth light</b>	130 μE
				<b>Toxicity</b>	Unproven
External Links					
<b>Genbank accession number</b>	-				
<b>References</b>	- B. Véron, J.-C. Dauguet and C. Billard. (1998) Sterolic biomarkers in marine phytoplankton. II. Free and conjugated sterols of 7 species used in mariculture. <i>Journal of Phycology</i> 34 (2): 273-279				