


## AC3 *Giraudyopsis stellifera*

Taxonomy		Image	Origin		
<b>Kingdom</b>	Chromista		<b>Climatic Zone</b>	Temperate	
<b>Division</b>	Heterokontophyta		<b>Habitat</b>	Coastal	
<b>Class</b>	Chrysomerophyceae		<b>Ocean</b>	North Atlantic	
<b>Order</b>	Chrysomeridales		<b>Sea</b>	English Channel	
<b>Family</b>	-		<b>Country</b>	France	
<b>Genus</b>	<i>Giraudyopsis</i>		<b>Region</b>	Brittany	
<b>Species</b>	stellifera		<b>Geolocation</b>		
<b>Authority(ies)</b>	Dangeard		<b>Latitude</b>	48.73	
			<b>Longitude</b>	-3.990000	
Identity		Morphology	Culture		
<b>Holotype</b>	No	<b>Length (µm)</b>	-	<b>Isolated by</b>	-
<b>Equivalent code</b>	-	<b>Width (µm)</b>	-	<b>Isolation date</b>	1969
<b>Deposited by</b>	Suzanne Loiseaux	<b>Diameter</b>	-	<b>Isolation site</b>	Roscoff
<b>Deposit date</b>	-	<b>Wall</b>	-	<b>Culture medium</b>	ES
<b>Identified by</b>	-	<b>Comments</b>	filaments, pseudokysts, zoospores with two unequal flagella slightly in ventral position (one smooth and one hairy)		
			<b>Growth temperature</b>	16 °C	
			<b>Growth light</b>	70 µE	
			<b>Toxicity</b>	Unproven	
External Links					
<b>Genbank accession number</b>	-				
<b>References</b>	<p>- Billard C. (1974). Microanatomie comparée de deux Chrysomméridacées. Individualisation des Sarcinochrysidales au sein des Chrysophycées. Soc. Phycol. de France, Bull. 19: 17-30</p> <p>- Billard C; J.C. Dauguet; D. Maume and Bert M. (1990). Sterols and chemotaxonomy of Marine Chrysophyceae. Botanica Marina 33: 225-228.</p>				