


AC30 *Hymenomonas globosa*

Taxonomy		Image		Origin	
Kingdom	Chromista			Climatic Zone	Temperate
Division	Haptophyta			Habitat	Coastal
Class	Prymnesiophyceae			Ocean	North Atlantic
Order	Coccolithales			Sea	English Channel
Family	<i>Hymenomonadaceae</i>			Country	France
Genus	<i>Hymenomonas</i>			Region	Basse-Normandie
Species	<i>globosa</i>			Geolocation	
Authority(ies)	(Magne) Gayral et Fresnel			Latitude	48.877
		Longitude	-1.831000		
Identity		Morphology		Culture	
Holotype	No	Length (μm)	-	Isolated by	C. Billard
Equivalent code	-	Width (μm)	-	Isolation date	1971
Deposited by	-	Diameter	-	Isolation site	Chausey (Îles)
Deposit date	-	Wall	-	Culture medium	ES
Identified by	-	Comments	Observed forms : palmelloid stage surrounded by coccoliths; zoide with coccoliths and two apical flagella	Growth temperature	16 °C
				Growth light	90 μE
				Toxicity	Unproven
External Links					
Genbank accession number	-				
References	- Fresnel J. (1994) A heteromorphic life cycle in two coastal coccolithophorids, <i>Hymenomonas lacuna</i> and <i>Hymenomonas coronata</i> (Prymnesiophyceae). Can. J. Bot. 72: 1455-1462				

AC30 *Hymenomonas globosa*

- Bittner L, Gobet A, Audic S, Romac S, Egge E.S., Santini S, Ogata H, Probert I, Edvardsen B and de Vargas C (2012) Diversity patterns of uncultured Haptophytes unravelled by pyrosequencing in Naples Bay. *Molecular Ecology* DOI: 10.1111/mec.12108