


AC32 *Pleurochrysis roscoffensis*

Taxonomy		Image		Origin	
Kingdom	Chromista			Climatic Zone	Temperate
Division	Haptophyta			Habitat	Coastal
Class	Prymnesiophyceae			Ocean	North Atlantic
Order	Coccolithales			Sea	English Channel
Family	<i>Pleurochrysidaceae</i>			Country	France
Genus	<i>Pleurochrysis</i>			Region	Brittany
Species	roscoffensis			Geolocation	
Authority(ies)	(Dangeard) Fresnel & Billard			Latitude	-
		Longitude	-		
Identity		Morphology		Culture	
Holotype	No	Length (µm)	-	Isolated by	J. Fresnel
Equivalent code	-	Width (µm)	-	Isolation date	1976
Deposited by	-	Diameter	-	Isolation site	Roscoff
Deposit date	-	Wall	-	Culture medium	f/2
Identified by	-	Comments	Small heterotroph contaminant present	Growth temperature	16 °C
				Growth light	160 µE
				Toxicity	Unproven
External Links					
Genbank accession number	-				
References	<p>- E. Plouguerné, C. Hellio, C. Cesconetto, M. Thabard, K. Mason, B. Véron, R. C. Pereira, B. A. P. da Gama (2010) Antifouling activity as a function of population variation in <i>Sargassum vulgare</i> from the littoral of Rio de Janeiro (Brazil). <i>Journal of Applied Phycology</i> 22 (6) : 717-724</p> <p>- Bittner L, Gobet A, Audic S, Romac S, Egge E.S., Santini S, Ogata H, Probert I, Edvardsen B and de Vargas C (2012) Diversity patterns of uncultured Haptophytes unravelled by pyrosequencing in Naples Bay. <i>Molecular Ecology</i> DOI: 10.1111/mec.12108</p>				