


# AC45 *Prymnesium parvum*

Taxonomy		Image		Origin	
<b>Kingdom</b>	Chromista			<b>Climatic Zone</b>	Temperate
<b>Division</b>	Haptophyta			<b>Habitat</b>	Coastal
<b>Class</b>	Prymnesiophyceae			<b>Ocean</b>	-
<b>Order</b>	Prymnesiales			<b>Sea</b>	-
<b>Family</b>	<i>Prymnesiaceae</i>			<b>Country</b>	France
<b>Genus</b>	<i>Prymnesium</i>			<b>Region</b>	Brittany
<b>Species</b>	parvum			<b>Geolocation</b>	
<b>Authority(ies)</b>	Carter			<b>Latitude</b>	48.674
				<b>Longitude</b>	-3.610000
Identity		Morphology		Culture	
<b>Holotype</b>	No	<b>Length (µm)</b>	-	<b>Isolated by</b>	C. Billard
<b>Equivalent code</b>	-	<b>Width (µm)</b>	-	<b>Isolation date</b>	1977
<b>Deposited by</b>	-	<b>Diameter</b>	-	<b>Isolation site</b>	Plestin-les-Grèves
<b>Deposit date</b>	-	<b>Wall</b>	-	<b>Culture medium</b>	ES
<b>Identified by</b>	C. Billard	<b>Comments</b>	Observed forms: solitary cells always flagellate.	<b>Growth temperature</b>	19 °C
				<b>Growth light</b>	100 µE
				<b>Toxicity</b>	Yes
External Links					
<b>Genbank accession number</b>	-				
<b>References</b>	Edvardsen et al. (2011). Ribosomal DNA phylogenies and a morphological revision provide the basis for a revised taxonomy of the Prymnesiales (Haptophyta). Eur. J. Phycol. 46 (3): 202-228				