


AC48 *Pleurochrysis roscoffensis*

Taxonomy		Image		Origin	
Kingdom	Chromista			Climatic Zone	Temperate
Division	Haptophyta			Habitat	Coastal
Class	Prymnesiophyceae			Ocean	North Atlantic
Order	Coccolithales			Sea	English Channel
Family	<i>Pleurochrysidaceae</i>			Country	France
Genus	<i>Pleurochrysis</i>			Region	Basse-Normandie
Species	roscoffensis			Geolocation	
Authority(ies)	-			Latitude	-
Identity		Morphology		Culture	
Holotype	No	Length (µm)	-	Isolated by	J. Fresnel
Equivalent code	-	Width (µm)	-	Isolation date	1979
Deposited by	-	Diameter	-	Isolation site	Courseulles-sur-mer
Deposit date	-	Wall	Coccosphere	Culture medium	f/2
Identified by	-	Comments	-	Growth temperature	16 °C
				Growth light	130 µE
				Toxicity	Unproven
External Links					
Genbank accession number	-				
References	- Gayral P, Fresnel J. (1983) Description, sexualité et cycle de développement d'une nouvelle coccolithophoracée (Prymnesiophyceae) : <i>Pleurochrysis pseudoroscoffensis</i> sp. nov. <i>Protistologica</i> XIX (2) : 245-261				

AC48 *Pleurochrysis roscoffensis*

- Bittner L, Gobet A, Audic S, Romac S, Egge E.S., Santini S, Ogata H, Probert I, Edvardsen B and de Vargas C (2012) Diversity patterns of uncultured Haptophytes unravelled by pyrosequencing in Naples Bay. *Molecular Ecology* DOI: 10.1111/mec.12108