

AC526 *Pleurochrysis* sp.

| Taxonomy | | Image | | Origin | |
|---------------------------------|---|--------------------|---|---------------------------|----------|
| Kingdom | Chromista | No image | | Climatic Zone | - |
| Division | Haptophyta | | | Habitat | - |
| Class | Prymnesiophyceae | | | Ocean | - |
| Order | Coccolithales | | | Sea | - |
| Family | <i>Pleurochrysidaceae</i> | | | Country | - |
| Genus | <i>Pleurochrysis</i> | | | Region | - |
| Species | sp. | | | Geolocation | |
| Authority(ies) | Pringsheim | | | Latitude | - |
| | | Longitude | - | | |
| Identity | | Morphology | | Culture | |
| Holotype | No | Length (µm) | - | Isolated by | - |
| Equivalent code | - | Width (µm) | - | Isolation date | - |
| Deposited by | - | Diameter | - | Isolation site | - |
| Deposit date | - | Wall | - | Culture medium | f/2 |
| Identified by | - | Comments | - | Growth temperature | 16 °C |
| | | | | Growth light | 50 µE |
| | | | | Toxicity | Unproven |
| External Links | | | | | |
| Genbank accession number | - | | | | |
| References | - Bittner L, Gobet A, Audic S, Romac S, Egge E.S., Santini S, Ogata H, Probert I, Edvardsen B and de Vargas C (2012) Diversity patterns of uncultured Haptophytes unravelled by pyrosequencing in Naples Bay. <i>Molecular Ecology</i> DOI: 10.1111/mec.12108 | | | | |