

AC570 *Prymnesium parvum*

| Taxonomy | | Image | | Origin | |
|---------------------------------|--|--------------------|---|---------------------------|----------|
| Kingdom | Chromista | No image | | Climatic Zone | - |
| Division | Haptophyta | | | Habitat | - |
| Class | Prymnesiophyceae | | | Ocean | - |
| Order | Prymnesiales | | | Sea | - |
| Family | <i>Prymnesiaceae</i> | | | Country | - |
| Genus | <i>Prymnesium</i> | | | Region | - |
| Species | parvum | | | Geolocation | |
| Authority(ies) | - | | | Latitude | - |
| | | Longitude | - | | |
| Identity | | Morphology | | Culture | |
| Holotype | No | Length (µm) | - | Isolated by | - |
| Equivalent code | - | Width (µm) | - | Isolation date | - |
| Deposited by | - | Diameter | - | Isolation site | - |
| Deposit date | - | Wall | - | Culture medium | K/2 |
| Identified by | - | Comments | - | Growth temperature | 16 °C |
| | | | | Growth light | 160 µE |
| | | | | Toxicity | Unproven |
| External Links | | | | | |
| Genbank accession number | - | | | | |
| References | Edwardsen et al. (2011). Ribosomal DNA phylogenies and a morphological revision provide the basis for a revised taxonomy of the Prymnesiales (Haptophyta). Eur. J. Phycol. 46 (3): 202-228 | | | | |