


AC58 *Hymenomonas coronata*

Taxonomy		Image	Origin		
Kingdom	Chromista		Climatic Zone	Subtropical	
Division	Haptophyta		Habitat	Inland water way	
Class	Prymnesiophyceae		Ocean	-	
Order	Coccolithales		Sea	-	
Family	<i>Hymenomonadaceae</i>		Country	États-Unis	
Genus	<i>Hymenomonas</i>		Region	Florida	
Species	coronata		Geolocation		
Authority(ies)	-		Latitude	-	
			Longitude	-	
Identity		Morphology	Culture		
Holotype	No	Length (μm)	-	Isolated by	J. Fresnel
Equivalent code	-	Width (μm)	-	Isolation date	08/1983
Deposited by	-	Diameter	-	Isolation site	Sarasota
Deposit date	-	Wall	-	Culture medium	ES
Identified by	-	Comments	-	Growth temperature	16 °C
				Growth light	90 μE
				Toxicity	Unproven
External Links					
Genbank accession number	-				
References	<p>- Fresnel J. (1994) A heteromorphic life cycle in two coastal coccolithophorids, <i>Hymenomonas lacuna</i> and <i>Hymenomonas coronata</i> (Prymnesiophyceae). <i>Can. J. Bot.</i> 72: 1455-1462</p> <p>- Bittner L, Gobet A, Audic S, Romac S, Egge E.S., Santini S, Ogata H, Probert I, Edvardsen B and de Vargas C (2012) Diversity patterns of uncultured Haptophytes unravelled by pyrosequencing in Naples Bay. <i>Molecular Ecology</i> DOI: 10.1111/mec.12108</p>				