


## AC609 *Chlamydomonas reinhardtii*

Taxonomy		Image		Origin	
<b>Kingdom</b>	Plantae			<b>Climatic Zone</b>	Temperate
<b>Division</b>	Chlorophyta			<b>Habitat</b>	Ground/soil
<b>Class</b>	Chlorophyceae			<b>Continent</b>	-
<b>Order</b>	Chlamydomonadales			<b>Country</b>	États-Unis
<b>Family</b>	<i>Chlamydomonadaceae</i>			<b>Region</b>	Massachusetts
<b>Genus</b>	<i>Chlamydomonas</i>			<b>Geolocation</b>	
<b>Species</b>	reinhardtii			<b>Latitude</b>	-
<b>Authority(ies)</b>	-			<b>Longitude</b>	-
Identity		Morphology		Culture	
<b>Holotype</b>	No	<b>Length (μm)</b>	-	<b>Isolated by</b>	G.M. Smith
<b>Equivalent code</b>	-	<b>Width (μm)</b>	-	<b>Isolation date</b>	1945
<b>Deposited by</b>	-	<b>Diameter</b>	6,10 μm	<b>Isolation site</b>	Amherst
<b>Deposit date</b>	-	<b>Wall</b>	-	<b>Culture medium</b>	L-C
<b>Identified by</b>	-	<b>Comments</b>	SAG: mating type dioecious Biovolume 0,121 pL	<b>Growth temperature</b>	16 °C
				<b>Growth light</b>	130 μE
				<b>Toxicity</b>	Unproven
External Links					
<b>Genbank accession number</b>	-				
<b>References</b>	-				