


AC738 *Porphyridium* sp.

| Taxonomy | | Image | Origin | | |
|---------------------------------|------------------------|--|----------------------|---------------------------|-----------------|
| Kingdom | Plantae |  | Climatic Zone | Mediterranean | |
| Division | Rhodophyta | | Habitat | Brackish | |
| Class | Porphyridiophyceae | | Ocean | - | |
| Order | Porphyridiales | | Sea | Mediterranean Sea | |
| Family | <i>Porphyridiaceae</i> | | Country | France | |
| Genus | <i>Porphyridium</i> | | Region | Languedoc-Roussillon | |
| Species | sp. | | Geolocation | | |
| Authority(ies) | - | | Latitude | - | |
| | | Longitude | - | | |
| Identity | | Morphology | Culture | | |
| Holotype | No | Length (μm) | - | Isolated by | G. Giraud |
| Equivalent code | - | Width (μm) | - | Isolation date | 1958 |
| Deposited by | - | Diameter | - | Isolation site | Banyuls-sur-Mer |
| Deposit date | 11 septembre 2008 | Wall | - | Culture medium | K/2 |
| Identified by | - | Comments | - | Growth temperature | 19 °C |
| | | | | Growth light | 100 μE |
| | | | | Toxicity | Unproven |
| External Links | | | | | |
| Genbank accession number | - | | | | |
| References | - | | | | |