


AC755 *Botryococcus braunii*

Taxonomy		Image		Origin	
Kingdom	Plantae			Climatic Zone	Temperate
Division	Chlorophyta			Habitat	Pool
Class	Trebouxiophyceae			Continent	Europe
Order	Trebouxiales			Country	France
Family	<i>Botryococcaceae</i>			Region	Bourgogne
Genus	<i>Botryococcus</i>			Geolocation	
Species	braunii			Latitude	-
Authority(ies)	Kützing			Longitude	-
Identity		Morphology		Culture	
Holotype	No	Length (µm)	10 µm	Isolated by	Metzger
Equivalent code	-	Width (µm)	5 µm	Isolation date	1981
Deposited by	Pierre Metzger, ENS Chimie, LCBOP, UMR CNRS 7618, 11 rue Pierre et Marie Curie 75231 PARIS Cedex 05, 0144276716, pierre-metzger@enscp.fr	Diameter	-	Isolation site	Lingoult-Morvan
Deposit date	28 octobre 2008	Wall	-	Culture medium	L-C
Identified by	-	Comments	-	Growth temperature	27 °C
				Growth light	-
				Toxicity	Unproven
External Links					
Genbank accession number	-				
References	- Senousy, H.H., Beakes, G.W., Hack, E. (2004). Phylogenetic placement of <i>Botryococcus braunii</i> . JPhycol. 40: 412-423)				

AC755 *Botryococcus braunii*

- Metzger, P., Berkaloff, C., Casadevall, E., Couté, A. (1985) Alkadiene- and botryococcene-producing races of wild strains of *Botryococcus braunii*. *Phytochemistry* 24: 2305-2312