


AC767 *Botryococcus braunii*

| Taxonomy | | Image | | Origin | |
|---------------------------------|---|--|------|---------------------------|-----------------|
| Kingdom | Plantae |  | | Climatic Zone | Subtropical |
| Division | Chlorophyta | | | Habitat | - |
| Class | Trebouxiophyceae | | | Continent | - |
| Order | Trebouxiales | | | Country | Thaïlande |
| Family | <i>Botryococcaceae</i> | | | Region | Songkla |
| Genus | <i>Botryococcus</i> | | | Geolocation | |
| Species | braunii | | | Latitude | - |
| Authority(ies) | Kützing | | | Longitude | - |
| | | | | | |
| Identity | | Morphology | | Culture | |
| Holotype | No | Length (µm) | 7 µm | Isolated by | Metzger |
| Equivalent code | - | Width (µm) | 3 µm | Isolation date | 1985 |
| Deposited by | - | Diameter | - | Isolation site | Songkla Nakarin |
| Deposit date | 28 octobre 2008 | Wall | - | Culture medium | L-C |
| Identified by | - | Comments | - | Growth temperature | 27 °C |
| | | | | Growth light | - |
| | | | | Toxicity | Unproven |
| External Links | | | | | |
| Genbank accession number | - | | | | |
| References | Senousy, H.H., Beakes, G.W., Hack, E. (2004). Phylogenetic placement of <i>Botryococcus braunii</i> . <i>JPhycol.</i> 40: 412-423), (Metzger, P., Casadevall, E. (1987) Lycopadiene, a tetraterpenoid hydrocarbon from new strains of <i>Botryococcus braunii</i> . <i>Tetrahedron Letters</i> 28: 3931-3934), | | | | |

AC767 *Botryococcus braunii*

(Metzger, P., Allard, B., Casadevall, E., Berkaloff, C., Couté, A. (1990). Structure and chemistry of a new chemical race of *Botryococcus braunii*. *JPhycol* 26: 258-266),

(Senousy, H.H. (2003) A molecular taxonomic and morphological study of the green algal genus *Botryococcus*. PhD Thesis, 248pp, University of Newcastle, UK.),

(Senousy, H.H., Beakes, G.W., Hack, E. (2004) Phylogenetic placement of *Botryococcus braunii* (Trebouxiophyceae) and *Botryococcus sudeticus* isolate UTEX 2629 (Chlorophyceae). *Jphycol* 40: 412-423